

REFERENCES

- [1]Tsuji H, Larson MG, Venditti Jr FJ, et al. Impact of reduced heart rate variability on risk for cardiac events: the Framingham Heart Study. *Circulation* 1996; 94(11): 2850–5.
- [2]Jerant A.F., Azari R., Nesbitt T.S.; Reducing the cost of frequent hospital admissions for congestive heart failure: a randomized trial of a home telecare intervention. *Med Care*. 39 2001:1234-1245.
- [3]SHUSTERMAN, V., andBARNEA, O. (1995): ‘Analysis of skin-temperature variability compared to variability of blood pressure and heart rate’,*IEEE Ann. Conf. Engineering Medicine Biology Society*, pp. 1027–1028.
- [4] C. Li, C. Zheng, and C. Tai, “Detection of eeg characteristic points using wavelet transforms,” *Biomed. Eng., IEEE Trans. on*, vol. 42, no. 1, pp. 21–28, 1995.
- [5]Dickstein K., Cohen-Solal A., Filippatos G.; ESC guidelines for the diagnosis and treatment of acute and chronic heart failure 2008: the Task Force for the Diagnosis and Treatment of Acute and Chronic Heart Failure 2008 of the European Society of Cardiology. Developed in collaboration with the Heart Failure Association of the ESC (HFA) and endorsed by the European Society of Intensive Care Medicine (ESICM). *Eur Heart J*. 29 2008:2388-2442.
- [6]Perini R, Orizio C, Milesi S, et al. Body position affects the power spectrum of heart rate variability during dynamic exercise. *Eur J Appl Physiol* 1993;